

Bohr-Sommerfeld tori and relative Poincaré series on a complex hyperbolic space

Tatyana Foth

February 8, 2008

Abstract

Automorphic forms on a bounded symmetric domain $D = G/K$ can be viewed as holomorphic sections of $L^{\otimes k}$, where L is a quantizing line bundle on a compact quotient of D and k is a positive integer.

Let Γ be a cocompact discrete subgroup of $SU(n, 1)$ which acts freely on $SU(n, 1)/U(n)$. We suggest a construction of relative Poincaré series associated to loxodromic elements of Γ . In complex dimension 2 we describe the Bohr-Sommerfeld tori in $\Gamma \backslash SU(n, 1)/U(n)$ associated to hyperbolic elements of Γ and prove that the relative Poincaré series associated to hyperbolic elements of Γ are not identically zero for large values of k .

1 Introduction

1.1 General definitions

We shall start with a brief review of the general concept of an automorphic form. Let G be a connected non-compact real semi-simple Lie group, K be a maximal compact subgroup of G , Γ be a discrete subgroup of G such that $\Gamma \backslash G$ has a finite volume. Let V be a finite-dimensional vector space, $\rho : K \rightarrow GL(V)$ be a representation of K . A smooth $Z(\mathfrak{g})$ -finite function $f : G \rightarrow V$ is called an **automorphic form on G for Γ** if

$$f(\gamma g k) = f(g) \rho(k) \quad (1)$$

for any $\gamma \in \Gamma$, $g \in G$, $k \in K$, and there are a positive constant C and a non-negative integer m such that

$$|f(g)| \leq C \|g\|^m \quad (2)$$

for any $g \in G$, here $|\cdot|$ is the norm corresponding to a $\rho(K)$ -invariant Hilbert structure on V , $\|g\| = \text{tr}(g^* g)$ taken in the adjoint representation of G . An automorphic form f is called a *cuspidal* form if $f \in L^\infty(\Gamma \backslash G)$.

The automorphy law (1) means geometrically that f defines a Γ -invariant section of the vector bundle $G \times_K V \rightarrow G/K$ associated to the principal bundle $G \rightarrow G/K$, here $G \times_K V = G \times V / \sim$, and the equivalence relation is given by the representation ρ : $(g, v) \sim (gk, v\rho(k))$.

The growth condition (2) is automatically satisfied with $m = 0$ in the case when $\Gamma \backslash G$ is compact and in this case any automorphic form is a cusp form.

Recall also that a function $f : G \rightarrow V$ is said to be **$Z(\mathfrak{g})$ -finite** if it is annihilated by an ideal I of $Z(\mathfrak{g})$ of a finite codimension, here $Z(\mathfrak{g})$ is the center of the universal enveloping algebra $U(\mathfrak{g})$.

$U(\mathfrak{g})$ can be identified with the algebra $D(G)$ of all left-invariant differential operators on G : to $Y \in \mathfrak{g}$ is associated a differential operator $Yf(g) = \frac{d}{dt}f(ge^{tY})|_{t=0}$, this establishes a linear map $\mathfrak{g} \rightarrow D(G)$ which extends to an isomorphism $U(\mathfrak{g}) \rightarrow D(G)$. $Z(\mathfrak{g})$ can be viewed as the subalgebra of all bi-invariant differential operators, it is isomorphic to a polynomial ring in l letters where l is the rank of G . A useful example to have in mind is $G = SL(2, R)$ and $\text{codim } I = 1$, then we have: $l = 1$, $Z(\mathfrak{g})$ is generated by the Casimir operator $\mathcal{C}$, and saying that a function f is $Z(\mathfrak{g})$ -finite is equivalent to stating that f is an eigenfunction of $\mathcal{C}$.

A well-known construction of an automorphic form on G is **Poincaré series**

$$\sum_{\gamma \in \Gamma} q(\gamma g),$$

where the function $q : G \rightarrow V$ is $Z(\mathfrak{g})$ -finite and K -finite on the right (i.e. the set of its right translates under elements of K is a finite-dimensional vector space), and $q \in L^1(G)$. One can also consider **relative Poincaré series**

$$\sum_{\gamma \in \Gamma_0 \backslash \Gamma} q(\gamma g),$$

where $q : G \rightarrow V$ is $Z(\mathfrak{g})$ -finite, K -finite on the right, Γ_0 -invariant, and $q \in L^1(\Gamma_0 \backslash G)$.

Let us explain now how to construct an automorphic form on G/K . An **automorphy factor** is a map $\mu : \Gamma \times G/K \rightarrow GL(V)$ such that $\mu(g_1 g_2, x) = \mu(g_1, g_2 x) \mu(g_2, x)$. It allows to define an **automorphic form on G/K** as a function $f : G/K \rightarrow V$ such that

$$f(\gamma x) \mu(\gamma, x) = f(x)$$

for any $\gamma \in \Gamma$, $x \in G/K$. Notice that then the function $F(g) = f(g(0)) \mu(g, 0)$, where $g \in G$, $x = g(0) \in G/K$, satisfies (1) with $\rho(k) = \mu(k, 0)$. Here 0 is the fixed point of K in G/K . If f is holomorphic then F is $Z(\mathfrak{g})$ -finite.

In particular, for a smooth function $q \in L^1(G/K)$ the Poincaré series on G/K is

$$\sum_{\gamma \in \Gamma} q(\gamma x) \mu(\gamma, x). \quad (3)$$

Similarly for a smooth Γ_0 -invariant function $q \in L^1(\Gamma_0 \backslash G/K)$ the relative Poincaré series is

$$\sum_{\gamma \in \Gamma_0 \backslash \Gamma} q(\gamma x) \mu(\gamma, x). \quad (4)$$

1.2 Automorphic forms on bounded symmetric domains and quantization

Consider a classical system (M, ω) , where M is a manifold, and ω is a symplectic form on M . The main problem of quantization is to associate a quantum system $(\mathcal{H}, \mathcal{O})$ to (M, ω) , where $\mathcal{H}$ is a Hilbert space and $\mathcal{O}$ consists of symmetric operators on $\mathcal{H}$.

The map $f \mapsto \hat{f}$, where $f \in C^\infty(M)$ and $\hat{f} \in \mathcal{O}$, should satisfy the following requirements: 1) it is $\mathbb{R}$ -linear, 2) if $f = \text{const}$ then $\hat{f}$ is the corresponding multiplication operator, 3) if $\{f_1, f_2\} = f_3$ then $\hat{f}_1 \hat{f}_2 - \hat{f}_2 \hat{f}_1 = -i\hbar \hat{f}_3$.

How do automorphic forms appear in the context of quantization ?

Suppose that M is a compact Kähler manifold which is a quotient of a bounded symmetric domain $D = G/K$ by the action of a discrete subgroup Γ , i.e. $M = \Gamma \backslash D$. Then the quantum phase space $\mathcal{H}$ consists of holomorphic automorphic forms on D for Γ . More precisely, let us consider the well-known quantization scheme for compact Kähler manifolds via Toeplitz operators (it is related to the standard scheme of geometric quantization with Kähler polarization). Then automorphic forms are holomorphic sections of $L^{\otimes k}$, where the canonical line bundle $L = \Lambda^n T^*M$ is the quantizing line bundle on M , here $n = \dim_{\mathbb{C}} M$ and k is a positive integer which determines the weight of an automorphic form, and $\hbar = \frac{1}{k}$.

We also notice that the automorphic forms (3) and (4) are sums of coherent states associated to holomorphic discrete series representations of G .

Let us describe all this in a bit more details. Let $D = G/K$ be a bounded symmetric domain, it is a Hermitian symmetric space of non-compact type (so the Riemannian metric on D is given by the real part of the hermitian form, and the symplectic form, which is a Kähler form in this case, is given by the imaginary part of the hermitian form, all these forms are G -invariant, of course). The irreducible Hermitian spaces of non-compact type are

- I) $SU(p, q)/S(U(p) \times U(q))$,
- II) $Sp(p, \mathbb{R})/U(p)$,
- III) $SO^*(2p)/U(p)$,
- IV) $SO_o(p, 2)/SO(p) \times SO(2)$

(and let us omit the case of an exceptional Lie algebra). So we have a metric

$$ds^2 = g_{ij} dz^i d\bar{z}^j, \quad (5)$$

the corresponding Kähler form is $\omega = g_{ij} dz^i \wedge d\bar{z}^j = i\partial\bar{\partial} \ln K(z, z)$, where $\frac{K(z, w)}{K(w, z)}$ is the Bergman kernel of the domain D . Recall that $K(z, w) = \overline{K(w, z)}$ and

$$K(\gamma z, \gamma w) = [\det J(\gamma, z)]^{-1} [\det \bar{J}(\gamma, w)]^{-1} K(z, w).$$

The Poisson bracket is

$$\{f, g\} = ig^{j\bar{l}} \left(\frac{\partial f}{\partial z^j} \frac{\partial g}{\partial \bar{z}^l} - \frac{\partial g}{\partial \bar{z}^j} \frac{\partial f}{\partial z^l} \right).$$

The **quantizing line bundle** $L \rightarrow M = \Gamma \backslash D$ can be defined as a line bundle such that the curvature of its natural connection is the Kähler form

ω on M . Denoting the canonical line bundle by L we see that the potential 1-form corresponding to the natural connection on L is $\Theta = i\partial \ln(s, s) = -i\partial \ln K(z, z)$, hence the curvature $d\Theta = -i\partial\bar{\partial} \ln K(z, z) = \omega$ and this is indeed a quantizing line bundle for M .

A holomorphic function $f : D \rightarrow \mathbb{C}$ is called an **automorphic form of weight k** if

$$f(\gamma z)[\det J(\gamma, z)]^k = f(z) \quad (6)$$

for any $z \in D$, $\gamma \in \Gamma$; here $J(\gamma, z)$ is the Jacobi matrix of transformation γ at point z . In the context of 1.1 the automorphy factor $\mu(\gamma, z) = [\det J(\gamma, z)]^k$. It is immediately clear that automorphic forms of weight k form the complex inner product space $H^0(M, L^{\otimes k})$ of holomorphic sections of $L^{\otimes k}$.

Now we consider a family of maps p_k , here k is a positive integer, such that $p_k(f) = T_f^{(k)}$, where f belongs to the Poisson algebra of smooth real-valued functions on M and $T_f^{(k)}$ is the Toeplitz operator on $H^0(M, L^{\otimes k})$ obtained from multiplication operator $M_f^{(k)}(g) = fg$ on $L^2(M, L^{\otimes k})$ by the orthogonal compression to the closed subspace $H^0(M, L^{\otimes k})$, i.e. $T_f^{(k)} = \Pi^{(k)} \circ M_f^{(k)} \circ \Pi^{(k)}$, where $\Pi^{(k)}$ is the orthogonal projection from $L^2(M, L^{\otimes k})$ to $H^0(M, L^{\otimes k})$.

In the Berezin scheme of quantization for each $\hbar = \frac{1}{k}$ we consider the space $\mathcal{F}_\hbar$ of functions holomorphic in D and satisfying (6) with the scalar product defined by

$$(f, g) = \text{const}(\hbar) \int_M f(z) \bar{g}(z) [K(z, z)]^{-\frac{1}{\hbar}} d\mu(z),$$

where $d\mu(z) = \omega^n$ is the G -invariant volume form on D corresponding to the metric (5). It is clear that $\mathcal{F}_\hbar$ is naturally identified with $H^0(M, L^{\otimes k})$. For the sake of completeness let us also explain briefly how the operator $\hat{A}$ corresponding a classical observable $A = A(z)$, is defined. First, we consider an analytic continuation $A(z, w)$ of the function $A(z)$ to $D \times D$. The covariant symbol $A(z, z)$ of $\hat{A}$ is defined as the diagonal value of the function

$$A(z, w) = \frac{\int_M \hat{A}[(K(u, w))^{\frac{1}{\hbar}}](K(z, u))^{\frac{1}{\hbar}} d\mu(u)}{\int_M (K(u, w))^{\frac{1}{\hbar}} (K(z, u))^{\frac{1}{\hbar}} d\mu(u)},$$

and

$$(\hat{A}f)(z) = \text{const}(\hbar) \int_M A(z, w) f(w) [K(z, w)]^{\frac{1}{\hbar}} [K(w, w)]^{-\frac{1}{\hbar}} d\mu(w).$$

So we end up with the algebra $A_\hbar$ of covariant symbols of bounded operators acting in $\mathcal{F}_\hbar$. The $*$ -product in $A_\hbar$ is given by

$$A_1 * A_2(z, z) = \text{const}(\hbar) \int_M A_1(z, w) A_2(w, z) \left(\frac{K(z, w) K(w, z)}{K(z, z) K(w, w)} \right)^{\frac{1}{\hbar}} d\mu(w).$$

In conclusion let us discuss the Poincaré series (3) and (4). Consider the unitary representation of G in $\mathcal{F}_\hbar$ given by the operators

$$[\pi^k(g)(f)](z) = [\det J(g^{-1}, z)]^k f(g^{-1}z).$$

It can be regarded as a subrepresentation of the left regular representation of G in $L^2(\Gamma \backslash G)$. Fix $f \in \mathcal{F}_h$, then the set $\{\pi^k(g)(f) | g \in G\}$ is called a *system of coherent states* (strictly speaking, we should regard two coherent states $\pi^k(g_1)(f)$ and $\pi^k(g_2)(f)$ as equivalent if $\pi^k(g_1)(f) = e^{i\alpha} \pi^k(g_2)(f)$). Now it is clear that (3) and (4) are just sums of coherent states corresponding to $f = q$ and the representation described above.

1.3 Comments on the subject of the present paper

In [9] and in the present paper we consider holomorphic automorphic forms on $D = \mathbb{H}_{\mathbb{C}}^n = SU(n, 1)/U(n)$. In [9] we construct a set of relative Poincaré series generating the graded algebra of $\mathbb{C}$ -valued cusp forms on a finite volume quotient of D . In the present paper we regard holomorphic $\mathbb{C}$ -valued automorphic forms on $\mathbb{H}_{\mathbb{C}}^n$ as holomorphic sections of the line bundle $L^{\otimes k} \rightarrow \Gamma \backslash \mathbb{H}_{\mathbb{C}}^n$, where L is a quantizing line bundle on $\Gamma \backslash \mathbb{H}_{\mathbb{C}}^n$, k is an integer, and Γ is a discrete cocompact subgroup of $SU(n, 1)$. We construct relative Poincaré series associated to loxodromic elements of Γ and we address an interesting problem which is not resolved for Poincaré series in general: is it true that these series are not identically zero? We restrict ourselves to the case of complex dimension 2 and answer “yes” to this question going through the following steps: 1) to each hyperbolic element of Γ we associate a certain Legendrian submanifold of the unit circle bundle in L^* such that the corresponding Lagrangian submanifold of $\Gamma \backslash SU(2, 1)/U(2)$ satisfies a Bohr-Sommerfeld condition, 2) following the method of [6] we conclude that the relative Poincaré series associated to hyperbolic elements are not zero for large values of k (i.e. in semi-classical limit $\hbar = \frac{1}{k} \rightarrow 0$).

2 Preliminaries

2.1 Complex hyperbolic space

Consider the complex hyperbolic space

$$\mathbb{H}_{\mathbb{C}}^n = SU(n, 1)/S(U(n) \times U(1)) = \mathbb{P}(\{z \in \mathbb{C}^{n+1} \mid \langle z, z \rangle < 0\}) \approx B^n,$$

here B^n is the open unit ball in $\mathbb{C}^n$, $\langle \cdot, \cdot \rangle$ is the Hermitian product on

$$\mathbb{C}^{n+1} \text{ given by } \langle z, w \rangle = z_1 \bar{w}_1 + \dots + z_n \bar{w}_n - z_{n+1} \bar{w}_{n+1} \text{ for } z = \begin{pmatrix} z_1 \\ \dots \\ z_{n+1} \end{pmatrix} \in \mathbb{C}^{n+1},$$

$$w = \begin{pmatrix} w_1 \\ \dots \\ w_{n+1} \end{pmatrix} \in \mathbb{C}^{n+1}.$$

A non-zero vector $z \in \mathbb{C}^{n+1}$ is called *negative (null, positive)* if the value of $\langle z, z \rangle$ is negative (null, positive).

$$\text{For } z, w \in B^n \text{ the corresponding vectors in } \mathbb{C}^{n+1} \text{ are } z = \begin{pmatrix} z_1 \\ \dots \\ z_n \\ 1 \end{pmatrix} \text{ and}$$

$$w = \begin{pmatrix} w_1 \\ \vdots \\ w_n \\ 1 \end{pmatrix}, \text{ and } \langle z, w \rangle = z_1 \bar{w}_1 + \dots + z_n \bar{w}_n - 1.$$

The group of biholomorphic automorphisms of $\mathbb{H}_{\mathbb{C}}^n$ is $PU(n, 1) = SU(n, 1)/\text{center}$. The group $SU(n, 1)$ acts on B^n by fractional-linear transformations: for

$$\gamma = \begin{pmatrix} a_{11} & \dots & a_{1n} & b_1 \\ \vdots & \vdots & \vdots & \vdots \\ a_{n1} & \dots & a_{nn} & b_n \\ c_1 & \dots & c_n & d \end{pmatrix}$$

we have:

$$\gamma z = \gamma(z_1, \dots, z_n) = \left(\frac{a_{11}z_1 + \dots + a_{1n}z_n + b_1}{c_1z_1 + \dots + c_nz_n + d}, \dots, \frac{a_{n1}z_1 + \dots + a_{nn}z_n + b_n}{c_1z_1 + \dots + c_nz_n + d} \right).$$

Notice that $\det J(\gamma, z) = (c_1z_1 + \dots + c_nz_n + d)^{-(n+1)}$, here $J(\gamma, z)$ denotes the Jacobi matrix of transformation γ at point $z \in B^n$.

An automorphism is called *loxodromic* if it has no fixed points in B^n and fixes two points in ∂B^n . Notice that the fixed points of the automorphisms correspond to the eigenvectors of the corresponding matrices in $U(n, 1)$. A loxodromic automorphism is called *hyperbolic* if it has a lift to $U(n, 1)$ all of whose eigenvalues are real.

A loxodromic element $\gamma_0 \in SU(n, 1)$ has $n - 1$ positive eigenvectors and two null eigenvectors.

Let $v_1, \dots, v_{n-1}$ be the positive eigenvectors of γ_0 , $\tau_1, \dots, \tau_{n-1}$ - the corresponding eigenvalues. Then $|\tau_j| = 1$, $1 \leq j \leq n - 1$.

Let X, Y be the null eigenvectors of γ_0 . Then the corresponding eigenvalues are λ and $\bar{\lambda}^{-1}$ for some $\lambda \in \mathbb{C}$, $|\lambda| > 1$.

A loxodromic transformation can always be represented by a matrix in $U(n, 1)$ with eigenvalues $\tau_1, \dots, \tau_{n-1}, \lambda, \bar{\lambda}^{-1}$ where $\lambda \in \mathbb{R}$, $|\lambda| > 1$.

The geodesic connecting X and Y (it is an arc of a circle orthogonal to ∂B^n or a diameter) is γ_0 -invariant and is called the *axis* of γ_0 . The complex line containing X and Y (*complex geodesic*) is γ_0 -invariant too.

2.2 Automorphic forms and geometry of the quotient

Let Γ be a discrete cocompact subgroup of $SU(n, 1)$ such that the quotient $X := \Gamma \backslash B^n$ is smooth.

A holomorphic function $f : B^n \rightarrow \mathbb{C}$ satisfying the automorphy law

$$f(\gamma z)(\det J(\gamma, z))^k = f(z) \quad (7)$$

for any $\gamma \in \Gamma$ is a cusp form of weight $(n + 1)k$ for Γ . The corresponding automorphic form on $SU(n, 1)$ is given by $F(g) = f(g(0))\zeta^k$, where $\zeta = \det J(g, 0)$, and the automorphy law on the group is

$$F(\gamma g) = F(g)$$

for any $\gamma \in \Gamma$. Notice that $\gamma : \zeta \rightarrow \zeta \det J(\gamma, z)$ for any $\gamma \in SU(n, 1)$.

We shall denote the space of cusp forms of weight $(n+1)k$ for Γ on B^n by $S_{(n+1)k}(\Gamma)$ and the corresponding space of cusp forms on $SU(n, 1)$ by $\tilde{S}_{(n+1)k}(\Gamma)$. The inner product on $S_{(n+1)k}(\Gamma)$ and $\tilde{S}_{(n+1)k}(\Gamma)$ is given by

$$(f, g) = (f(z)\zeta^k, g(z)\zeta^k) = i^n \int_{\Gamma \backslash B^n} f\bar{g}(-\langle z, z \rangle)^{(n+1)k} \frac{dz_1 \wedge d\bar{z}_1 \wedge \dots \wedge dz_n \wedge d\bar{z}_n}{(-\langle z, z \rangle)^{n+1}}.$$

Given a subgroup Γ_0 of Γ and a holomorphic function $q(z) \in L^2(\Gamma \backslash B^n)$ satisfying (7) for all $\gamma \in \Gamma_0$, the *relative Poincaré series* for Γ_0 is defined as

$$\Theta(z) = \sum_{\gamma \in \Gamma_0 \backslash \Gamma} q(\gamma z) (\det J(\gamma, z))^k,$$

this series converges absolutely and uniformly on compact sets and belongs to the space $S_{(n+1)k}(\Gamma)$.

The Bergman kernel for the domain B^n is $K(z, w) = \frac{1}{(-\langle z, w \rangle)^{n+1}}$ and the Kähler form on X is

$$\Phi_\kappa = \frac{2\kappa i}{\langle z, z \rangle^2} (\langle z, z \rangle \sum_{j=1}^n dz_j \wedge d\bar{z}_j - \langle dz, z \rangle \wedge \langle z, dz \rangle)$$

where κ is a positive constant, it is an $SU(n, 1)$ -invariant Kähler form on B^n .

Remark 2.1 We will set $\kappa = \frac{n+1}{2}$. Then the holomorphic sectional curvature on B^n is $-\frac{2}{\kappa} = -\frac{4}{n+1}$ and the sectional curvature is pinched between $-\frac{4}{n+1}$ and $-\frac{1}{n+1}$ ([10] II.2.2, III.1.5).

Consider the canonical line bundle $L := \bigwedge^n T^*X$ and the dual bundle $L^* = \bigwedge^n TX$. The potential 1-form θ on L^* is characterized by

$$\nabla s = -i\theta s,$$

where ∇ is a connection on L^* and s is the unit section. The potential 1-form corresponding to the natural connection on L^* is

$$\theta = i\partial \ln(s, s) = i\partial \ln((-\langle z, z \rangle)^{n+1}).$$

Here $d = \partial + \bar{\partial}$. The curvature on L^* is $d\theta = -\Phi_{\frac{n+1}{2}}$, hence L is the natural quantizing line bundle for X .

Let

$$E_k = H^0(X, L^{\otimes k})$$

be the complex inner-product space of holomorphic sections of the k -th tensor power of L . Consider the unit circle bundle $P \subset L^*$, a point of P can be described as (z, ζ) , where $z \in B^n$ and ζ is the coordinate on the fiber, $|\zeta| = (-\langle z, z \rangle)^{\frac{n+1}{2}}$. We have:

$$E_k = \tilde{S}_{(n+1)k}(\Gamma) = \{f(z)\zeta^k \mid (z, \zeta) \in P, f \text{ is holomorphic on } B^n,$$

$$f(\gamma z)(\det J(\gamma, z))^k = f(z) \text{ for any } \gamma \in \Gamma\}.$$

Denote also $\tilde{L} = \bigwedge^n T^*B^n$, $\tilde{P}$ - the unit circle bundle in $\tilde{L}^*$.

The connection form $\alpha : TP \rightarrow \mathbb{R}$ on P is

$$\alpha = \theta + i \frac{d\zeta}{\zeta}.$$

It serves as a contact form on $\tilde{P}$ and P .

A Lagrangian submanifold $\Lambda_0 \subset X$ satisfies a *Bohr-Sommerfeld condition* if

$$\frac{k}{2\pi} \int_C \theta \in \mathbb{Z}$$

for any closed curve $C \subset \Lambda_0$. The constant $\frac{1}{k}$ plays role of the Planck constant.

The unit disk bundle W in L^* is a compact, strictly pseudoconvex domain with smooth boundary. Let us consider the *Hardy space* of P : $E \subset L^2(P)$ and the *Szëgo projector*

$$\Pi : L^2(P) \rightarrow E$$

given by the orthogonal projection of L^2 onto E . We identify:

$$E = \oplus_{k=0}^{\infty} E_k.$$

Let $\tilde{E}$ be the Hardy space of $\tilde{P}$, then $\tilde{E} = \oplus_{k=0}^{\infty} \tilde{E}_k$ where

$$\tilde{E}_k = \{f(z)\zeta^k \mid (z, \zeta) \in \tilde{P}, f \text{ is holomorphic on } B^n,$$

$$i^n \int_{B^n} |f(z)|^2 (-\langle z, z \rangle)^{(n+1)k} \frac{dz_1 \wedge d\bar{z}_1 \wedge \dots \wedge dz_n \wedge d\bar{z}_n}{(-\langle z, z \rangle)^{n+1}} < \infty\}.$$

We shall denote the corresponding orthogonal projection by

$$\tilde{\Pi} : L^2(\tilde{P}) \rightarrow \tilde{E}.$$

3 Construction of relative Poincaré series associated to certain loxodromic elements of a discrete cocompact subgroup of $SU(n,1)$

Consider a loxodromic automorphism of B^n , represented it by a matrix $\gamma_0 \in U(n, 1)$ with eigenvalues $\tau_1, \dots, \tau_{n-1}, \lambda, \lambda^{-1}$, $|\tau_j| = 1$, $j = 1, \dots, n-1$, $\lambda \in \mathbb{R}$, $|\lambda| > 1$, denote the corresponding eigenvectors by $v_1, \dots, v_{n-1}, X, Y$ ($v_1, \dots, v_{n-1}$ are positive, X, Y are null). Notice that if each τ_j is a root of 1 then some power of γ_0 is a hyperbolic element.

Assumption 3.1 Assume that 1 is among the eigenvalues of γ_0 .

Remark 3.2 If $g \in U(n, 1)$ is hyperbolic then g^2 is a hyperbolic element of $SU(n, 1)$ which satisfies Assumption 3.1 and has the same eigenvectors as g .

Generalizing the construction suggested in [9], for any collection, w.l.o.g. $v_1, \dots, v_m$, $m \leq n-1$, of positive eigenvectors corresponding to eigenvalue 1 we construct a relative Poincaré series

$$\Theta_{\gamma_0, l, k} = \sum_{\gamma \in \Gamma_0 \setminus \Gamma} q_l(\gamma z) (\det J(\gamma, z))^{2k} \in S_{2(n+1)k},$$

where $\Gamma_0 = \langle \gamma_0 \rangle$,

$$q_l(z) = \frac{\langle z, v_1 \rangle^{l_1} \dots \langle z, v_m \rangle^{l_m}}{(\langle z, X \rangle \langle z, Y \rangle)^{(n+1)k + \frac{l_1 + \dots + l_m}{2}}},$$

$l_1, \dots, l_m$ are positive integers such that $l_1 + \dots + l_m$ is even, $l = (l_1, \dots, l_m)$. The series converges absolutely in B^n and uniformly on the compact sets by the Theorem .9 for $k \geq 1$ (see the Appendix).

In dimension 2 the loxodromic elements of Γ satisfying Assumption 3.1 are exactly the hyperbolic elements of Γ . The relative Poincaré series associated to a hyperbolic element $\gamma_0 \in \Gamma$ is

$$\Theta_{\gamma_0, l, k} = \sum_{\gamma \in \Gamma_0 \setminus \Gamma} q_l(\gamma z) (\det J(\gamma, z))^{2k} \in S_{6k},$$

where $\Gamma_0 = \langle \gamma_0 \rangle$,

$$q_l(z) = \frac{\langle z, v \rangle^{2l}}{(\langle z, X \rangle \langle z, Y \rangle)^{3k+l}},$$

and l is a positive integer.

Remark 3.3 Let γ_1 and γ_2 be hyperbolic elements of Γ . If $\gamma_1 = \gamma_2^N$ for a positive integer N , then $\Theta_{\gamma_1, l, k} = \Theta_{\gamma_2, l, k}$. If γ_1 and γ_2 are conjugate in Γ then $\Theta_{\gamma_1, l, k} = \Theta_{\gamma_2, l, k}$.

4 Bohr-Sommerfeld tori

Consider a hyperbolic element $\gamma_0 \in \Gamma$, denote its null eigenvectors by X , Y , denote its positive eigenvector by v , then the corresponding eigenvalues are λ , λ^{-1} , 1, for $\lambda \in \mathbb{R}$, $|\lambda| > 1$.

We choose v so that

$$A = [v \quad \frac{X}{\langle X, Y \rangle} + \frac{Y}{2} \quad \frac{X}{\langle X, Y \rangle} - \frac{Y}{2}]$$

belongs to $SU(2, 1)$.

The transformation A^{-1} maps the complex line containing X and Y to the complex line $\{z_1 = 0\}$ and maps the geodesic connecting X and Y to the geodesic $\tilde{C}$ connecting $(0, -1)$ and $(0, 1)$. The following loxodromic element of $SU(2, 1)$ preserves $\tilde{C}$ and the line $\{z_1 = 0\}$:

$$\gamma := A^{-1} \gamma_0 A = \begin{pmatrix} 1 & 0 & 0 \\ 0 & a & b \\ 0 & b & a \end{pmatrix},$$

$$a = \frac{\lambda^2 + 1}{2\lambda}, \quad b = \frac{\lambda^2 - 1}{2\lambda}, \quad a^2 - b^2 = 1.$$

Denote

$$w = (w_1, w_2) = A^{-1}z, \quad w_1 = A^{-1}z_1, \quad w_2 = A^{-1}z_2,$$

and apply change of variables

$$w_2 = \frac{re^{i\phi} - i}{re^{i\phi} + i}, \quad w_1 = \sqrt{1 - w_2\bar{w}_2}Re^{i\Theta},$$

$$0 < \phi < \pi, \quad 0 < r < +\infty, \quad 0 < R < 1, \quad 0 \leq \Theta < 2\pi.$$

Proposition 4.1 *Any 2-cylinder $C_{\phi,R} = \{\phi = \text{const}, R = \text{const}\}$ is γ -invariant.*

Remark 4.2 *The coordinates (r, Θ) on $C_{\phi,R}$ are the “radial” and the “angular” coordinates respectively.*

Proof.

$$w_2 \rightarrow \frac{aw_2 + b}{bw_2 + a} = \frac{a \frac{re^{i\phi} - i}{re^{i\phi} + i} + b}{b \frac{re^{i\phi} - i}{re^{i\phi} + i} + a} = \frac{r(a+b)e^{i\phi} - i(a-b)}{r(a+b)e^{i\phi} + i(a-b)} = \frac{r \frac{a+b}{a-b} e^{i\phi} - i}{r \frac{a+b}{a-b} e^{i\phi} + i},$$

so

$$r \rightarrow r \frac{a+b}{a-b}, \quad \phi \rightarrow \phi,$$

also

$$\frac{|w_1|}{\sqrt{1 - w_2\bar{w}_2}} \rightarrow \frac{\left| \frac{w_1}{bw_2 + a} \right|}{\sqrt{1 - \left| \frac{aw_2 + b}{bw_2 + a} \right|^2}} = \frac{|w_1|}{\sqrt{1 - w_2\bar{w}_2}},$$

so $R \rightarrow R$.

○

For a positive integer l consider the following submanifold of $\tilde{P}$:

$$\tilde{T}(l) := \{(w, (-\langle w, w \rangle)^{\frac{3}{2}} e^{i\psi}) \mid w \in \{\phi = \frac{\pi}{2}, R = \sqrt{\frac{l}{3k+l}}\}, \psi = -\frac{l}{k}\Theta\}.$$

Denote $T(l) := \langle \gamma \rangle \setminus \tilde{T}(l)$, $\tilde{\Lambda}(l) := A\tilde{T}(l)$ and $\Gamma_0 := \langle \gamma_0 \rangle$.

Proposition 4.3 $\Lambda(l) := AT(l) = \Gamma_0 \setminus \tilde{\Lambda}(l)$ is a compact Legendrian submanifold of P .

Proof. $T(l)$ and $\Lambda(l)$ are compact submanifolds of P .

Let us prove that $\Lambda(l)$ is Legendrian. Submanifolds $T(l)$ and $\Lambda(l)$ have dimension 2. The restriction of α onto $\tilde{T}(l)$ is

$$\begin{aligned} -3i \frac{\langle dw, w \rangle}{\langle w, w \rangle} + i \frac{d\zeta}{\zeta} &= -3i \frac{\langle dw, w \rangle}{\langle w, w \rangle} - d\psi + i \frac{d(-\langle w, w \rangle)^{\frac{3}{2}}}{(-\langle w, w \rangle)^{\frac{3}{2}}} = \\ -3i \frac{\langle dw, w \rangle}{\langle w, w \rangle} - d\psi - \frac{3}{2} i \frac{\langle dw, w \rangle + \langle w, dw \rangle}{(-\langle w, w \rangle)^{\frac{3}{2}}} (-\langle w, w \rangle)^{\frac{1}{2}} &= \\ -3i \frac{\langle dw, w \rangle}{\langle w, w \rangle} - d\psi + \frac{3}{2} i \frac{\langle dw, w \rangle}{\langle w, w \rangle} + \frac{3}{2} i \frac{\langle w, dw \rangle}{\langle w, w \rangle} &= \end{aligned}$$

$$\begin{aligned}
& -\frac{3}{2}i \frac{\bar{w}_1 dw_1 + \bar{w}_2 dw_2}{w_1 \bar{w}_1 + w_2 \bar{w}_2 - 1} + \frac{3}{2}i \frac{w_1 d\bar{w}_1 + w_2 d\bar{w}_2}{w_1 \bar{w}_1 + w_2 \bar{w}_2 - 1} - d\psi = \\
& -\frac{3}{2}i \frac{(1 - w_2 \bar{w}_2) R^2 e^{-i\Theta} i e^{i\Theta} d\Theta + \sqrt{1 - w_2 \bar{w}_2} R^2 e^{-i\Theta} e^{i\Theta} d\sqrt{1 - w_2 \bar{w}_2} + \bar{w}_2 dw_2}{(1 - w_2 \bar{w}_2) R^2 + w_2 \bar{w}_2 - 1} + \\
& \frac{3}{2}i \frac{-(1 - w_2 \bar{w}_2) R^2 e^{-i\Theta} i e^{i\Theta} d\Theta + \sqrt{1 - w_2 \bar{w}_2} R^2 e^{-i\Theta} e^{i\Theta} d\sqrt{1 - w_2 \bar{w}_2} + w_2 d\bar{w}_2}{(1 - w_2 \bar{w}_2) R^2 + w_2 \bar{w}_2 - 1} - d\psi = \\
& -\frac{3}{2}i \frac{(1 - w_2^2) R^2 id\Theta + \sqrt{1 - w_2^2} R^2 d\sqrt{1 - w_2^2} + w_2 dw_2}{(1 - w_2^2)(R^2 - 1)} + \\
& \frac{3}{2}i \frac{-(1 - w_2^2) R^2 id\Theta + \sqrt{1 - w_2^2} R^2 d\sqrt{1 - w_2^2} + w_2 dw_2}{(1 - w_2^2)(R^2 - 1)} - d\psi = \\
& -3i \frac{(1 - w_2^2) R^2 id\Theta}{(1 - w_2^2)(R^2 - 1)} - d\psi = 3 \frac{R^2 d\Theta}{R^2 - 1} - d\psi = 3 \frac{\frac{l}{3k+l} d\Theta}{\frac{l}{3k+l} - 1} + \frac{l}{k} d\Theta = 0.
\end{aligned}$$

We showed that $T(l)$ is a Legendrian submanifold of P . To prove that $AT(l)$ is Legendrian too it is enough to show that α is $SU(2, 1)$ -invariant.

Let $M \in SU(2, 1)$,

$$M : (z, \zeta) \rightarrow (Mz, \zeta \det J(M, z)) = (Mz, \zeta c^3),$$

where $c = c(z) = (m_{31}z_1 + m_{32}z_2 + 1)^{-1}$.

$$\alpha = i \frac{d\zeta}{\zeta} - 3i \frac{\langle dz, z \rangle}{\langle z, z \rangle} = i \frac{d\zeta}{\zeta} - 3i \partial \ln(-\langle z, z \rangle).$$

We have:

$$\begin{aligned}
& i \frac{d(c^3 \zeta)}{c^3 \zeta} - 3i \partial \ln(-\langle Mz, Mz \rangle) = i \frac{c^3 d\zeta + 3c^2 \zeta dc}{c^3 \zeta} - 3i \partial \ln(-\langle z, z \rangle c \bar{c}) = \\
& i \frac{d\zeta}{\zeta} + 3i \frac{dc}{c} - 3i \partial \ln(-\langle z, z \rangle) - 3i \partial \ln(c \bar{c}) = i \frac{d\zeta}{\zeta} + 3i \frac{\partial c}{c} - 3i \partial \ln(-\langle z, z \rangle) - 3i \frac{\partial c}{c} = \\
& i \frac{d\zeta}{\zeta} - 3i \partial \ln(-\langle z, z \rangle).
\end{aligned}$$

○

The natural projection $\Lambda_0(l)$ of $\Lambda(l)$ onto X is a compact Lagrangian submanifold of X .

Proposition 4.4 $\Lambda_0(l)$ satisfies a Bohr-Sommerfeld condition.

Proof. Let $\tilde{T}_0(l)$ be the natural projection of $\tilde{T}(l)$ onto B^2 , and let $T_0(l)$ be the natural projection of $T(l)$ onto X , $AT_0(l) = \Lambda_0(l)$. If $C \subset \Lambda_0(l)$ is a closed curve then $A^{-1}C \subset T_0(l)$ is also closed. Let $z \in \Lambda_0(l)$, $w \in T_0(l)$, $c = c(w) = (a_{31}w_1 + a_{32}w_2 + a_{33})^{-1}$, we have:

$$\begin{aligned}
\int_C \theta &= -3i \int_C \partial \ln(-\langle z, z \rangle) = -3i \int_{A^{-1}C} \partial \ln(-\langle Aw, Aw \rangle) = \\
& -3i \int_{A^{-1}C} \partial \ln(-\langle w, w \rangle c \bar{c}) =
\end{aligned}$$

$$\begin{aligned}
-3i \int_{A^{-1}C} \partial \ln(-\langle w, w \rangle) + \frac{\bar{c} \partial c}{c \bar{c}} &= -3i \int_{A^{-1}C} \partial \ln(-\langle w, w \rangle) + \partial \ln c = \\
-3i \int_{A^{-1}C} \partial \ln(-\langle w, w \rangle) + d \ln c &= -3i \int_{A^{-1}C} \partial \ln(-\langle w, w \rangle),
\end{aligned}$$

so $\int_C \theta$ is A^{-1} -invariant (in fact $SU(2, 1)$ -invariant) and it is enough to prove that $T_0(l)$ satisfies the Bohr-Sommerfeld condition. The restriction of the potential 1-form onto $\tilde{T}_0(l)$ is

$$\begin{aligned}
& -3i \frac{\bar{w}_1 dw_1 + \bar{w}_2 dw_2}{w_1 \bar{w}_1 + w_2 \bar{w}_2 - 1} = \\
& -3i \frac{(1 - w_2 \bar{w}_2) R^2 e^{-i\Theta} i e^{i\Theta} d\Theta + \sqrt{1 - w_2 \bar{w}_2} R^2 e^{-i\Theta} e^{i\Theta} d\sqrt{1 - w_2 \bar{w}_2} + \bar{w}_2 dw_2}{(1 - w_2 \bar{w}_2) R^2 + w_2 \bar{w}_2 - 1} = \\
& -3i \frac{(1 - w_2^2) R^2 i d\Theta + \sqrt{1 - w_2^2} R^2 d\sqrt{1 - w_2^2} + w_2 dw_2}{(1 - w_2^2)(R^2 - 1)} = \\
& -3i \frac{(1 - w_2^2) R^2 i d\Theta + \sqrt{1 - w_2^2} R^2 \frac{-2w_2 dw_2}{2\sqrt{1 - w_2^2}} + w_2 dw_2}{(1 - w_2^2)(R^2 - 1)} = \\
& -3i \left(\frac{R^2 i}{R^2 - 1} d\Theta - \frac{w_2 dw_2}{1 - w_2^2} \right) = \\
& 3 \frac{R^2}{R^2 - 1} d\Theta - 3i \frac{1}{2} d \ln(1 - w_2^2) = -\frac{l}{k} d\Theta - \frac{3}{2} i d \ln(1 - w_2^2),
\end{aligned}$$

then

$$\begin{aligned}
& \frac{3k}{2\pi} \int_{A^{-1}C} \left(-\frac{l}{k} d\Theta - \frac{3}{2} i d \ln(1 - w_2^2) \right) = \\
& \frac{-3l}{2\pi} \int_{A^{-1}C} d\Theta = \frac{-3l}{2\pi} 2\pi m = -3lm \in \mathbb{Z}.
\end{aligned}$$

○

So the torus $\Lambda_0(l)$ is a Lagrangian submanifold satisfying the Bohr-Sommerfeld condition.

Proposition 4.5 *The orthogonal projection of the delta function at $(w, \eta) \in \tilde{P}$ into $\tilde{E}_k$ is the function*

$$\Psi_{(w, \eta)}(z, \zeta) := \tilde{\Pi}_k(\delta_{(w, \eta)}) = \frac{(3k-1)(3k-2)}{4\pi^2} \frac{\zeta^k \bar{\eta}^k}{\langle z, w \rangle^{3k}}.$$

Remark 4.6 *The orthogonal projection of the delta function at $(w, \eta) \in \tilde{P}$ into $\tilde{E}_k$ is the coherent state in $\tilde{E}_k$ associated to the point $(w, \eta) \in \tilde{P}$, by definition $g\Psi_{(w, \eta)} = \Psi_{g(w, \eta)}$ for $g \in SU(2, 1)$.*

Proof. The fact that $\Psi_{(w, \eta)} = \tilde{\Pi}_k(\delta_{(w, \eta)})$ is equivalent to the reproducing property:

$$F(w, \eta) = \int_{\tilde{P}} \bar{\Psi}_{(w, \eta)}(z, \zeta) F(z, \zeta) dV \wedge d\psi$$

for all $F \in \tilde{E}_k$. Given any orthonormal basis $\{F_{l,k}\}$ for $\tilde{E}_k$, we can write the reproducing kernel as the series $\Psi_{(w, \eta)}(z, \zeta) = \sum_l \bar{F}_{l,k}(w, \eta) F_{l,k}(z, \zeta)$ which converges absolutely and uniformly on compact sets.

Using the basis

$$F_{l,m,k}(z, \zeta) = \frac{1}{2\pi} \sqrt{\frac{(3k+l+m-1)!}{l!m!(3k-3)!}} z_1^l z_2^m \zeta^k,$$

which is orthonormal with respect to the inner product

$$(f(z)\zeta^k, g(z)\zeta^k) = i^2 \int_{B^2} f\bar{g}(-\langle z, z \rangle)^{3k-3} dz_1 \wedge d\bar{z}_1 \wedge dz_2 \wedge d\bar{z}_2,$$

we obtain:

$$\begin{aligned} \Psi_{(w,\eta)}(z, \zeta) &= \sum_{l,m} \bar{F}_{l,m,k}(w, \eta) F_{l,m,k}(z, \zeta) = \\ &= \sum_{l,m} \frac{1}{(2\pi)^2} \frac{(3k+l+m-1)!}{l!m!(3k-3)!} \bar{w}_1^l \bar{w}_2^m z_1^l z_2^m \zeta^k \bar{\eta}^k = \\ &= \frac{\zeta^k \bar{\eta}^k}{4\pi^2 (3k-3)!} \sum_{l,m} \frac{(3k+l+m-1)!}{l!m!} (\bar{w}_1 z_1)^l (\bar{w}_2 z_2)^m. \end{aligned}$$

To calculate

$$\sum_{l,m} \frac{(3k+l+m-1)!}{l!m!} x^l y^m = \sum_m \frac{y^m}{m!} \sum_l \frac{(3k+l+m-1)!}{l!} x^l$$

let us find first $\sum_l \frac{(N+l)!}{l!} t^l$. Integrating once we get $\sum_{l=0}^{\infty} \frac{(N+l)!}{(l+1)!} t^{l+1}$; integrating twice we get $\sum_{l=0}^{\infty} \frac{(N+l)!}{(l+2)!} t^{l+2}$; integrating N times we get:

$$\sum_{l=0}^{\infty} \frac{(N+l)!}{(l+N)!} t^{l+N} = \sum_{l=0}^{\infty} t^{l+N} = \frac{t^N}{1-t},$$

differentiating this expression N times we get:

$$\begin{aligned} \sum_l \frac{(N+l)!}{l!} t^l &= \left(\frac{t^N}{1-t} \right)^{(N)} = \left(\frac{t^N - 1 + 1}{1-t} \right)^{(N)} = \\ &= \left(\frac{(t-1)(t^{N-1} + t^{N-2} + \dots + t + 1) + 1}{1-t} \right)^{(N)} = \left(\frac{1}{1-t} \right)^{(N)} = \frac{N!}{(1-t)^{N+1}}, \end{aligned}$$

hence

$$\begin{aligned} \sum_m \frac{y^m}{m!} \sum_l \frac{(3k+l+m-1)!}{l!} x^l &= \sum_m \frac{y^m}{m!} \frac{(3k+m-1)!}{(1-x)^{3k+m}} = \\ &= \frac{1}{(1-x)^{3k}} \sum_m \frac{(3k+m-1)!}{m!} \left(\frac{y}{1-x} \right)^m = \frac{1}{(1-x)^{3k}} \frac{(3k-1)!}{\left(1 - \frac{y}{1-x}\right)^{3k}} = \frac{(3k-1)!}{(1-x-y)^{3k}}, \end{aligned}$$

so

$$\Psi_{(w,\eta)}(z, \zeta) = \frac{\zeta^k \bar{\eta}^k}{4\pi^2 (3k-3)!} \frac{(3k-1)!}{(1 - \bar{w}_1 z_1 - \bar{w}_2 z_2)^{3k}} = \frac{(3k-1)(3k-2)}{4\pi^2} \zeta^k \bar{\eta}^k \frac{1}{(-\langle z, w \rangle)^{3k}}.$$

○

For weight $6k$ we have:

$$\Psi_{(u,\eta)}(z, \zeta) = \tilde{\Pi}_{2k}(\delta_{(u,\eta)}) = \frac{(6k-1)(6k-2)}{4\pi^2} \frac{\zeta^{2k} \bar{\eta}^{2k}}{\langle z, u \rangle^{6k}}.$$

We omit the weight in the notation $\Psi_{(u,\eta)}(z, \zeta)$ but further exposition will be for weight $6k$ so this will not lead to any confusion.

To get the orthogonal projection of the delta function at $[(u, \eta)] \in P = \Gamma \setminus \tilde{P}$ (by $[(u, \eta)]$ we denote the equivalence class of (u, η)) into E_{2k} we average over the action of Γ :

$$\Pi_{2k}(\delta_{[(u,\eta)]}) = \sum_{g \in \Gamma} g \Psi_{(u,\eta)}. \quad (8)$$

The function $\Psi_{(u,\eta)}$ belongs to $\tilde{E}_{2k}$, hence the series (8) converges absolutely and uniformly on compact sets by Theorem 9.1 [4].

Following the method of [6], to the submanifold $\Lambda(l) \subset P$ with a half-form ν we associate a function

$$\begin{aligned} \Phi &:= \int_{\Lambda(l)} \Pi_{2k}(\delta_{[(u,\eta)]}) \nu = \sum_{g \in \Gamma} \int_{\Lambda(l)} g \Psi_{(u,\eta)} \nu = \\ &= \sum_{g \in \Gamma/\Gamma_0} \sum_{m=-\infty}^{+\infty} \int_{\Lambda(l)} g \gamma_0^m \Psi_{(u,\eta)}(z, \zeta) \nu = \sum_{g \in \Gamma/\Gamma_0} \sum_{m=-\infty}^{+\infty} \int_{\Lambda(l)} g \Psi_{\gamma_0^m(u,\eta)}(z, \zeta) \nu = \\ &= \sum_{g \in \Gamma/\Gamma_0} g \int_{\tilde{\Lambda}(l)} \Psi_{(u,\eta)}(z, \zeta) \nu = \sum_{g^{-1} \in \Gamma_0 \setminus \Gamma} g^{-1} \int_{\tilde{\Lambda}(l)} \Psi_{(u,\eta)}(z, \zeta) \nu. \end{aligned}$$

Proposition 4.7

$$\int_{\tilde{\Lambda}(l)} \Psi_{(u,\eta)}(z, \zeta) \nu = C \frac{\langle z, v \rangle^{2l}}{(\langle z, X \rangle \langle z, Y \rangle)^{3k+l}} \zeta^{2k},$$

where the constant C is given by

$$C = 2^{3k+l-2} \frac{i}{(2l)!} \frac{(6k+2l-1)!}{(6k-3)!} \frac{(3k)^{3k} l^l}{(3k+l)^{3k+l}} \langle Y, X \rangle^{3k+l} \sum_{j=0}^{3k+l-1} \frac{(-1)^j (3k+l-1)!}{j! j! (6k+2l-1-j)!}.$$

and we take

$$\nu = \frac{d(A^{-1}u_1)}{A^{-1}u_1} \wedge \frac{d(A^{-1}u_2)}{1 - (A^{-1}u_2)^2}.$$

Remark 4.8 The half-form ν on $\tilde{\Lambda}(l)$ is γ_0 -invariant and in properly chosen coordinates (r, Θ) on $\tilde{\Lambda}_0(l)$ it is expressed as $\nu = \frac{i}{2} d\Theta \wedge \frac{dr}{r}$.

Proof. Let $u \in \tilde{\Lambda}(l)$, $w = A^{-1}u \in \tilde{T}(l)$, then

$$\begin{aligned} \int_{\tilde{\Lambda}(l)} \Psi_{(u,\eta)}(z, \zeta) \nu &= \frac{(6k-1)(6k-2)}{4\pi^2} \int_{\tilde{\Lambda}(l)} \frac{\zeta^{2k} \bar{\eta}^{2k}}{\langle z, u \rangle^{6k}} \frac{d(A^{-1}u_1)}{A^{-1}u_1} \wedge \frac{d(A^{-1}u_2)}{1 - (A^{-1}u_2)^2} = \\ &= \frac{(6k-1)(6k-2)}{4\pi^2} \zeta^{2k} \int_{\tilde{\Lambda}(l)} \frac{\bar{\eta}^{2k}}{\langle z, u \rangle^{6k}} \frac{d(A^{-1}u_1)}{A^{-1}u_1} \wedge \frac{d(A^{-1}u_2)}{1 - (A^{-1}u_2)^2} = \end{aligned}$$

$$\begin{aligned} & \frac{(6k-1)(6k-2)}{4\pi^2} \zeta^{2k} \int_{\tilde{T}(l)} \frac{((- \langle w, w \rangle)^{\frac{3}{2}} e^{-i\psi} \det \bar{J}(A, w))^{2k}}{\langle z, Aw \rangle^{6k}} \frac{dw_1}{w_1} \wedge \frac{dw_2}{1-w_2^2} = \\ & \frac{(6k-1)(6k-2)}{4\pi^2} \zeta^{2k} \int_{\tilde{T}(l)} \frac{(- \langle w, w \rangle)^{3k} e^{-i2k\psi} (\det \bar{J}(A, w))^{2k}}{\langle A^{-1}z, w \rangle^{6k} (\det \bar{J}(A, w))^{2k}} (\det J(A^{-1}, z))^{2k} \frac{dw_1}{w_1} \wedge \frac{dw_2}{1-w_2^2}, \end{aligned}$$

let $A^{-1}z = \begin{pmatrix} v_1 \\ v_2 \\ 1 \end{pmatrix}$, then we get:

$$\frac{(6k-1)(6k-2)}{4\pi^2} \zeta^{2k} (\det J(A^{-1}, z))^{2k} \int_{\tilde{T}(l)} \frac{(- \langle w, w \rangle)^{3k} e^{-i2k\psi}}{(v_1 \bar{w}_1 + v_2 \bar{w}_2 - 1)^{6k}} \frac{dw_1}{w_1} \wedge \frac{dw_2}{1-w_2^2},$$

on $\tilde{T}(l)$

$$\begin{aligned} w_2 &= \frac{r-1}{r+1}, \quad w_1 = \sqrt{1-w_2^2} R e^{i\Theta} = \frac{2\sqrt{r}}{r+1} R e^{i\Theta}, \\ - \langle w, w \rangle &= (1-R^2)(1-w_2^2) = (1-R^2) \frac{4r}{(r+1)^2}, \end{aligned}$$

so we have:

$$\begin{aligned} & \frac{(6k-1)(6k-2)}{4\pi^2} \zeta^{2k} (\det J(A^{-1}, z))^{2k} (4(1-R^2))^{3k} \\ & \int_{\tilde{T}(l)} \frac{(\frac{r}{(r+1)^2})^{3k} e^{-i2k\psi}}{(v_1 \frac{2\sqrt{r}}{r+1} R e^{-i\Theta} + v_2 \frac{r-1}{r+1} - 1)^{6k}} \frac{i}{2} d\Theta \wedge \frac{dr}{r} = \\ & \frac{(6k-1)(6k-2)}{4\pi^2} \zeta^{2k} (\det J(A^{-1}, z))^{2k} (4(1-R^2))^{3k} \frac{i}{2} \\ & \int_0^\infty dr \int_0^{2\pi} d\Theta \frac{r^{3k-1} e^{i2l\Theta}}{(v_1 2\sqrt{r} R e^{-i\Theta} + v_2(r-1) - r - 1)^{6k}}. \end{aligned}$$

The integral

$$\int_{|w|=1} \frac{1}{(Aw+B)^{6k}} \frac{dw}{w^{2l+1}}, \quad \left| \frac{B}{A} \right| > 1$$

is equal to

$$\frac{2\pi i}{(2l)!} \frac{d^{2l}}{dw^{2l}} \frac{1}{(Aw+B)^{6k}} \Big|_{w=0} = \frac{2\pi i}{(2l)!} \frac{(6k+2l-1)!}{(6k-1)!} \frac{A^{2l}}{B^{6k+2l}},$$

Let $w = e^{-i\Theta}$, $A = v_1 2\sqrt{r}R$, $B = v_2(r-1) - r - 1$. Let us check that $|\frac{B}{A}| > 1$.

$$\begin{aligned} \left| \frac{v_2(r-1) - r - 1}{v_1 2\sqrt{r}R} \right| &= \left| \frac{v_2 w_2 - 1}{v_1 R \sqrt{1-w_2^2}} \right| > \frac{|v_2 w_2 - 1|}{\sqrt{1-v_2 \bar{v}_2} R \sqrt{1-w_2^2}} \geq \\ & \frac{|v_2 w_2 - 1|}{\sqrt{1-v_2 \bar{v}_2} \sqrt{1-w_2^2}} \geq 1 \end{aligned}$$

because

$$\begin{aligned} 0 \leq |v_2 - w_2|^2 &= (\bar{v}_2 - w_2)(v_2 - w_2) = v_2 \bar{v}_2 - \bar{v}_2 w_2 - w_2 v_2 + w_2^2 = \\ & -\bar{v}_2 w_2 - w_2 v_2 + v_2 \bar{v}_2 w_2^2 + 1 + v_2 \bar{v}_2 + w_2^2 - v_2 \bar{v}_2 w_2^2 - 1 = \end{aligned}$$

$$(\bar{v}_2 w_2 - 1)(v_2 w_2 - 1) - (1 - v_2 \bar{v}_2)(1 - w_2^2).$$

We get:

$$\begin{aligned} & \frac{(6k-1)(6k-2)}{4\pi^2} \zeta^{2k} (4(1-R^2))^{3k} \frac{1}{2} \frac{2\pi i}{(2l)!} \frac{(6k+2l-1)!}{(6k-1)!} (\det J(A^{-1}, z))^{2k} \\ & \int_0^\infty \frac{r^{3k-1+l} v_1^{2l} 2^{2l} R^{2l}}{(v_2(r-1) - r - 1)^{6k+2l}} dr = \\ & \frac{\pi i}{(2l)!} \frac{1}{4\pi^2} \zeta^{2k} (4(1-R^2))^{3k} \frac{(6k+2l-1)!}{(6k-3)!} \frac{v_1^{2l} 2^{2l} R^{2l}}{(v_2-1)^{6k+2l}} (\det J(A^{-1}, z))^{2k} \int_0^\infty \frac{r^{3k-1+l}}{(r - \frac{v_2+1}{v_2-1})^{6k+2l}} dr. \end{aligned}$$

The integral $\int_0^\infty \frac{r^{3k+l-1}}{(r-a)^{6k+2l}} dr = -\text{res}_{z=a} f(z)$ where $f(z) = \frac{z^{3k+l-1} \ln z}{(z-a)^{6k+2l}}$ ([22] 29.6). Let us calculate this residue:

$$\begin{aligned} \text{res}_{z=a} f(z) &= \frac{1}{(6k+2l-1)!} \frac{d^{6k+2l-1}}{dz^{6k+2l-1}} z^{3k+l-1} \ln z|_{z=a} = \\ & \frac{1}{(6k+2l-1)!} \sum_{j=0}^{6k+2l-1} C_{6k+2l-1}^j (z^{3k+l-1})^{(j)} (\ln z)^{(6k+2l-1-j)}|_{z=a} = \\ & \frac{1}{(6k+2l-1)!} \sum_{j=0}^{3k+l-1} C_{6k+2l-1}^j (z^{3k+l-1})^{(j)} \left(\frac{1}{z}\right)^{(6k+2l-2-j)}|_{z=a} = \\ & \frac{1}{(6k+2l-1)!} \sum_{j=0}^{3k+l-1} \frac{(6k+2l-1)!}{j!(6k+2l-1-j)!} \frac{(3k+l-1)!}{j!} z^{3k+l-1-j} \\ & \left. \frac{(6k+2l-2-j)!(-1)^{6k+2l-2-j}}{z^{6k+2l-1-j}} \right|_{z=a} = C_1 \frac{1}{a^{3k+l}}, \end{aligned}$$

where $C_1 = \sum_{j=0}^{3k+l-1} \frac{(-1)^j (3k+l-1)!}{j! j! (6k+2l-1-j)!}$. Hence

$$\int_0^\infty \frac{r^{3k+l-1}}{(r - \frac{v_2+1}{v_2-1})^{6k+2l}} dr = -C_1 \left(\frac{v_2-1}{v_2+1}\right)^{3k+l}$$

and we get:

$$\begin{aligned} & -\frac{\pi i}{(2l)!} \frac{1}{4\pi^2} \zeta^{2k} (4(1-R^2))^{3k} \frac{(6k+2l-1)!}{(6k-3)!} C_1 \frac{v_1^{2l} 2^{2l} R^{2l}}{(v_2-1)^{6k+2l}} (\det J(A^{-1}, z))^{2k} \left(\frac{v_2-1}{v_2+1}\right)^{3k+l} = \\ & \frac{i}{(2l)!} \frac{1}{4\pi} \zeta^{2k} (4(1-R^2))^{3k} \frac{(6k+2l-1)!}{(6k-3)!} C_1 2^{2l} R^{2l} (\det J(A^{-1}, z))^{2k} \frac{A^{-1} z_1^{2l}}{((A^{-1} z_2)^2 - 1)^{3k+l}} = \\ & 2^{6k+2l-2} \frac{i}{(2l)!} \frac{(6k+2l-1)!}{(6k-3)!} C_1 \left(\frac{3k}{3k+l}\right)^{3k} \left(\frac{l}{3k+l}\right)^l \left(\frac{1}{2} \langle Y, X \rangle\right)^{-3k-l} \frac{\langle z, v \rangle^{2l}}{(\langle z, X \rangle \langle z, Y \rangle)^{3k+l}} \zeta^{2k}. \end{aligned}$$

○

We got:

$$\Phi(z, \zeta) = C \zeta^{2k} \sum_{g \in \Gamma_0 \setminus \Gamma} q_l(gz) (\det J(g, z))^{2k} \in E_k,$$

where

$$q_l(z) = \frac{\langle z, v \rangle^{2l}}{(\langle z, X \rangle \langle z, Y \rangle)^{3k+l}}$$

and the relative Poincaré series associated to $\Lambda_0(l)$ is

$$\Theta_{\gamma_0, l, k}(z) := C \sum_{g \in \Gamma_0 \setminus \Gamma} q_l(gz) (\det J(g, z))^{2k}.$$

From the results of [6] (Theorem 3.2, Corollary 3.3) it follows that $\Theta_{\gamma_0, l, k}$ is non-vanishing for sufficiently large values of k .

APPENDIX.

We shall prove the following theorem modifying the proof of convergence of Poincaré series contained in [4] and [3].

Theorem .9 *Let φ be a function on $G = SU(n, 1)$. Assume that*

- 1) φ is Z -finite
 - 2) $\varphi \in L^1(\Gamma_0 \setminus G)$,
 - 3) φ is K -finite on the right
- Let $p_\varphi(x) = \sum_{\gamma \in \Gamma_0 \setminus \Gamma} \varphi(\gamma x)$.*

Then p_φ converges absolutely and uniformly on compact sets.

Proof.

By Lemma 9.2 [4] there exists $\alpha \in C_c^\infty(G)$ satisfying $\alpha(k^{-1}xk) = \alpha(x)$, $k \in K$, $x \in G$, such that $\varphi = \varphi * \alpha$. such that $U^{-1} = U$, the closure of U is compact, and $U \supset \text{supp } \alpha$. We have:

$$\varphi(\gamma x) = (\varphi * \alpha)(\gamma x) = \int_G \varphi(\gamma xy) \alpha(y^{-1}) dy = \int_U \varphi(\gamma xy) \alpha(y^{-1}) dy,$$

hence

$$|\varphi(\gamma x)| \leq \|\alpha\|_\infty \int_U |\varphi(\gamma xy)| dy = \|\alpha\|_\infty \int_{xU} |\varphi(\gamma y)| dy$$

Here $\|\alpha\|_\infty = \sup_{y \in U} |\alpha(y)|$.

Fix a compact subset C of G . We want to prove absolute and uniform convergence on C . The closure of CU is compact. CU is covered by N copies of a "fundamental domain" of Γ in G , N is a positive integer (because Γ is discrete). Denote these domains by $F_1, \dots, F_N$. By a "fundamental domain" of Γ in G I mean a connected set of representatives of $\Gamma \setminus G$.

Let $x \in C$. Then

$$|\varphi(\gamma x)| \leq \|\alpha\|_\infty \int_{xU} |\varphi(\gamma y)| dy \leq \|\alpha\|_\infty \int_{CU} |\varphi(\gamma y)| dy$$

and we get

$$\sum_{\gamma \in \Gamma_0 \setminus \Gamma} \|\alpha\|_\infty \int_{CU} |\varphi(\gamma y)| dy = \|\alpha\|_\infty \sum_{\gamma \in \Gamma_0 \setminus \Gamma} \int_{CU} |\varphi(\gamma y)| dy \leq$$

$$\begin{aligned}
\|\alpha\|_\infty \sum_{\gamma \in \Gamma_0 \setminus \Gamma} \left(\int_{F_1} |\varphi(\gamma y)| dy + \dots + \int_{F_N} |\varphi(\gamma y)| dy \right) = \\
\|\alpha\|_\infty \left(\sum_{\gamma \in \Gamma_0 \setminus \Gamma} \int_{F_1} |\varphi(\gamma y)| dy + \dots + \sum_{\gamma \in \Gamma_0 \setminus \Gamma} \int_{F_N} |\varphi(\gamma y)| dy \right) = \\
N \|\alpha\|_\infty \int_{\Gamma_0 \setminus G} |\varphi(y)| dy < \infty.
\end{aligned}$$

So we proved that

$$|\varphi(\gamma x)| \leq c_\gamma := \|\alpha\|_\infty \int_{CU} |\varphi(\gamma y)| dy$$

and that the numerical series $\sum_{\gamma \in \Gamma_0 \setminus \Gamma} c_\gamma$ converges, hence by Weierstrass theorem the series $\sum_{\gamma \in \Gamma_0 \setminus \Gamma} \varphi(\gamma x)$ converges absolutely and uniformly on C . Q.E.D.

References

- [1] F. Berezin, *General concept of quantization*, Comm. Math. Phys. **40** (1975), 153-174.
- [2] M. Bordemann, E. Meinrenken, and M. Schlichenmaier, *Toeplitz quantization of Kähler manifolds and $gl(N)$, $N \rightarrow \infty$ limits*, Comm. Math. Phys. **165** (1994), no. 2, 281-296.
- [3] A. Borel, *Automorphic forms on $SL_2(\mathbb{R})$* , Cambridge University Press, 1997.
- [4] A. Borel, *Introduction to automorphic forms*, Proc. Symp. Pure Math. **9** (1966), 199-210.
- [5] D. Borthwick, A. Lesniewski, and H. Upmeyer, *Non-perturbative deformation quantization of Cartan domains*, J. Funct. Anal. **113** (1993), 153-176.
- [6] D. Borthwick, T. Paul, and A. Uribe, *Legendrian distributions with applications to relative Poincaré series*, Invent. math. **122** (1995), 359-402.
- [7] L. Boutet de Monvel and V. Guillemin, *The spectral theory of Toeplitz operators*, Princeton University Press, 1981.
- [8] D. Epstein, *Complex hyperbolic geometry*, in *Analytical and geometric aspects of hyperbolic space*, London Math. Soc. Lecture Note Ser., 111, Cambridge Univ. Press, 1987.
- [9] T. Foth and S. Katok, *Spanning sets for automorphic forms and dynamics of the frame flow on complex hyperbolic spaces*, Submitted.
- [10] W. Goldman, *Complex hyperbolic geometry*, Oxford University Press, 1999.
- [11] V. Guillemin and S. Sternberg, *Geometric asymptotics*, Math. Surveys no. 14, AMS, 1977.

- [12] N. Hurt, *Geometric quantization in action*, D. Reidel Publ. Co., 1983.
- [13] A. Kirillov, *Elements of the theory of representations*, Springer-Verlag, 1976.
- [14] S. Klimek and A. Lesniewski, *Quantum Riemann surfaces I. The unit disc*, Comm. Math. Phys. **146** (1992), 103-122.
- [15] S. Klimek and A. Lesniewski, *Quantum Riemann surfaces II. The discrete series*, Lett. Math. Phys. **24** (1992), 125-139.
- [16] J. Kollar, *Shafarevich maps and automorphic forms*, Princeton University Press, 1995.
- [17] S. Krantz, *Function theory of several complex variables*, Wadsworth & Brooks/Cole, 1992.
- [18] A. Perelomov, *Generalized coherent states and applications*, Springer-Verlag, 1986.
- [19] J. Rawnsley, *Quantization on Kähler manifolds*, in *Differential geometric methods in theoretical physics*, Lect. Notes in Phys. **375** (1991), 155-161.
- [20] J. Rawnsley, *Deformation quantization of Kähler manifolds*, in *Symplectic geometry and mathematical physics*, Prog. in Math. **99** (1991), 366-373.
- [21] Y. Tai and H. Resnikoff, *On the structure of a graded ring of automorphic forms on the 2-dimensional complex ball, I and II*, Math. Ann. **238**(1978), 97-117; Math. Ann. **258**(1982), 367-382.
- [22] Yu. Sidorov, M. Fedoryuk, and M. Shabunin, *Lectures on the theory of functions of a complex variable*, Mir Publishers, Moscow, 1985.
- [23] J. Sniatycki, *Geometric quantization and quantum mechanics*, Springer-Verlag, 1980.
- [24] D. Zhelobenko and A. Shtern, *Representations of Lie groups (Russian)*, "Nauka", Moscow, 1983.
- [25] N. Woodhouse, *Geometric quantization*, Oxford University Press, 1980.